

表 1103 脳梗塞の既往 PTH値(pg/mL)別（腹膜透析患者）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	33	7	40			40
(%)	(82.5)	(17.5)	(100.0)			
20～	42	3	45			45
(%)	(93.3)	(6.7)	(100.0)			
35～	48	6	54		1	55
(%)	(88.9)	(11.1)	(100.0)			
50～	64	12	76		1	77
(%)	(84.2)	(15.8)	(100.0)			
75～	61	7	68		1	69
(%)	(89.7)	(10.3)	(100.0)			
100～	63	8	71			71
(%)	(88.7)	(11.3)	(100.0)			
125～	40	5	45			45
(%)	(88.9)	(11.1)	(100.0)			
150～	1		1			1
(%)	(100.0)		(100.0)			
151～	38	4	42		1	43
(%)	(90.5)	(9.5)	(100.0)			
180～	27	5	32		2	34
(%)	(84.4)	(15.6)	(100.0)			
210～	23	4	27			27
(%)	(85.2)	(14.8)	(100.0)			
240～	21	2	23			23
(%)	(91.3)	(8.7)	(100.0)			
300～	17	2	19			19
(%)	(89.5)	(10.5)	(100.0)			
350～	7	1	8			8
(%)	(87.5)	(12.5)	(100.0)			
400～	8	3	11			11
(%)	(72.7)	(27.3)	(100.0)			
500～	9	2	11			11
(%)	(81.8)	(18.2)	(100.0)			
合計	502	71	573		6	579
(%)	(87.6)	(12.4)	(100.0)			
記載なし	94	16	110		30	140
(%)	(85.5)	(14.5)	(100.0)			
総計	596	87	683		36	719
(%)	(87.3)	(12.7)	(100.0)			
平均	129.85	152.56	132.66		122.33	132.56
標準偏差	119.23	188.98	129.91		67.35	129.39

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査