

表 1117 四肢切断の有無 原疾患別（透析患者全体）

原疾患	切断なし	切断あり	合計	不明	記載なし	総計
慢性糸球体腎炎	73,061	1,004	74,065	82	22,823	96,970
(%)	(98.6)	(1.4)	(100.0)			
慢性腎盂腎炎	2,323	42	2,365	1	676	3,042
(%)	(98.2)	(1.8)	(100.0)			
急速進行性糸球体腎炎	1,885	24	1,909	1	584	2,494
(%)	(98.7)	(1.3)	(100.0)			
妊娠腎/妊娠中毒症	1,251	15	1,266	2	346	1,614
(%)	(98.8)	(1.2)	(100.0)			
その他の分類不能の腎炎	1,029	6	1,035	1	337	1,373
(%)	(99.4)	(0.6)	(100.0)			
多発性嚢胞腎	8,435	56	8,491	13	2,502	11,006
(%)	(99.3)	(0.7)	(100.0)			
腎硬化症	21,884	299	22,183	58	6,057	28,298
(%)	(98.7)	(1.3)	(100.0)			
悪性高血圧	1,924	28	1,952	4	617	2,573
(%)	(98.6)	(1.4)	(100.0)			
糖尿病性腎症	84,134	6,649	90,783	108	27,190	118,081
(%)	(92.7)	(7.3)	(100.0)			
SLE腎炎	1,710	33	1,743	5	488	2,236
(%)	(98.1)	(1.9)	(100.0)			
アミロイド腎	333	2	335	1	115	451
(%)	(99.4)	(0.6)	(100.0)			
痛風腎	869	16	885	3	225	1,113
(%)	(98.2)	(1.8)	(100.0)			
先天性代謝異常による腎不全	200	4	204	1	67	272
(%)	(98.0)	(2.0)	(100.0)			
腎・尿路結核	150	2	152		54	206
(%)	(98.7)	(1.3)	(100.0)			
腎・尿路結石	437	6	443		138	581
(%)	(98.6)	(1.4)	(100.0)			
腎・尿路腫瘍	656	5	661	1	224	886
(%)	(99.2)	(0.8)	(100.0)			
閉塞性尿路障害	554	21	575		148	723
(%)	(96.3)	(3.7)	(100.0)			
骨髄腫	210	1	211		65	276
(%)	(99.5)	(0.5)	(100.0)			
腎形成不全	505	8	513		136	649
(%)	(98.4)	(1.6)	(100.0)			
不明	20,053	384	20,437	29	7,078	27,544
(%)	(98.1)	(1.9)	(100.0)			
再導入	1,620	45	1,665		498	2,163
(%)	(97.3)	(2.7)	(100.0)			
その他	5,584	137	5,721	10	1,826	7,557
(%)	(97.6)	(2.4)	(100.0)			
合計	228,807	8,787	237,594	320	72,194	310,108
(%)	(96.3)	(3.7)	(100.0)			
記載なし						
(%)						
総計	228,807	8,787	237,594	320	72,194	310,108
(%)	(96.3)	(3.7)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査