

表 1134 四肢切断の有無 PTH値(pg/mL)別（透析患者全体）PTH測定法:whole-PTH

PTH値	切断なし	切断あり	合計	不明	記載なし	総計
20未満	1,543	65	1,608		286	1,894
(%)	(96.0)	(4.0)	(100.0)			
20～	1,924	72	1,996		335	2,331
(%)	(96.4)	(3.6)	(100.0)			
35～	1,686	65	1,751		292	2,043
(%)	(96.3)	(3.7)	(100.0)			
50～	2,559	98	2,657		496	3,153
(%)	(96.3)	(3.7)	(100.0)			
75～	2,204	91	2,295		453	2,748
(%)	(96.0)	(4.0)	(100.0)			
100～	1,854	51	1,905		393	2,298
(%)	(97.3)	(2.7)	(100.0)			
125～	1,206	38	1,244		295	1,539
(%)	(96.9)	(3.1)	(100.0)			
150～	38	2	40		8	48
(%)	(95.0)	(5.0)	(100.0)			
151～	1,016	38	1,054		209	1,263
(%)	(96.4)	(3.6)	(100.0)			
180～	710	22	732		143	875
(%)	(97.0)	(3.0)	(100.0)			
210～	510	16	526		102	628
(%)	(97.0)	(3.0)	(100.0)			
240～	547	12	559		115	674
(%)	(97.9)	(2.1)	(100.0)			
300～	230	4	234		50	284
(%)	(98.3)	(1.7)	(100.0)			
350～	145	9	154		25	179
(%)	(94.2)	(5.8)	(100.0)			
400～	169	1	170		32	202
(%)	(99.4)	(0.6)	(100.0)			
500～	201	9	210		37	247
(%)	(95.7)	(4.3)	(100.0)			
合計	16,542	593	17,135		3,271	20,406
(%)	(96.5)	(3.5)	(100.0)			
記載なし	371	23	394		263	657
(%)	(94.2)	(5.8)	(100.0)			
総計	16,913	616	17,529		3,534	21,063
(%)	(96.5)	(3.5)	(100.0)			
平均	107.63	98.37	107.31		111.65	108.01
標準偏差	113.18	99.75	112.75		120.36	114.01

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査