

表 1135 四肢切断の有無 透析前ヘモグロビン濃度(g/dL)別（透析患者全体）

透析前ヘモグロビン濃度	切断なし	切断あり	合計	不明	記載なし	総計
5未満	256	13	269		34	303
(%)	(95.2)	(4.8)	(100.0)			
5～	355	15	370		59	429
(%)	(95.9)	(4.1)	(100.0)			
6～	920	42	962	3	195	1,160
(%)	(95.6)	(4.4)	(100.0)			
7～	3,510	193	3,703	7	725	4,435
(%)	(94.8)	(5.2)	(100.0)			
8～	11,614	609	12,223	29	2,302	14,554
(%)	(95.0)	(5.0)	(100.0)			
9～	36,809	1,517	38,326	60	7,603	45,989
(%)	(96.0)	(4.0)	(100.0)			
10～	75,577	2,541	78,118	89	15,705	93,912
(%)	(96.7)	(3.3)	(100.0)			
11～	62,940	2,130	65,070	79	13,024	78,173
(%)	(96.7)	(3.3)	(100.0)			
12～	23,174	906	24,080	32	4,742	28,854
(%)	(96.2)	(3.8)	(100.0)			
13～	5,842	271	6,113	4	1,191	7,308
(%)	(95.6)	(4.4)	(100.0)			
14～	2,153	119	2,272	2	399	2,673
(%)	(94.8)	(5.2)	(100.0)			
合計	223,150	8,356	231,506	305	45,979	277,790
(%)	(96.4)	(3.6)	(100.0)			
記載なし	5,657	431	6,088	15	26,215	32,318
(%)	(92.9)	(7.1)	(100.0)			
総計	228,807	8,787	237,594	320	72,194	310,108
(%)	(96.3)	(3.7)	(100.0)			
平均	10.71	10.65	10.71	10.49	10.71	10.71
標準偏差	1.28	1.42	1.29	1.31	1.26	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査