

表 1145 四肢切断の有無 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	切断なし	切断あり	合計	不明	記載なし	総計
慢性糸球体腎炎	55,617	777	56,394	70	12,113	68,577
(%)	(98.6)	(1.4)	(100.0)			
慢性腎盂腎炎	1,785	32	1,817	1	347	2,165
(%)	(98.2)	(1.8)	(100.0)			
急速進行性糸球体腎炎	1,514	21	1,535	1	298	1,834
(%)	(98.6)	(1.4)	(100.0)			
妊娠腎/妊娠中毒症	879	14	893	1	170	1,064
(%)	(98.4)	(1.6)	(100.0)			
その他の分類不能の腎炎	764	3	767		182	949
(%)	(99.6)	(0.4)	(100.0)			
多発性嚢胞腎	6,750	46	6,796	10	1,392	8,198
(%)	(99.3)	(0.7)	(100.0)			
腎硬化症	17,849	241	18,090	45	3,325	21,460
(%)	(98.7)	(1.3)	(100.0)			
悪性高血圧	1,523	24	1,547	3	331	1,881
(%)	(98.4)	(1.6)	(100.0)			
糖尿病性腎症	69,523	5,413	74,936	93	15,478	90,507
(%)	(92.8)	(7.2)	(100.0)			
SLE腎炎	1,276	29	1,305	2	243	1,550
(%)	(97.8)	(2.2)	(100.0)			
アミロイド腎	264	1	265	1	59	325
(%)	(99.6)	(0.4)	(100.0)			
痛風腎	693	12	705	1	122	828
(%)	(98.3)	(1.7)	(100.0)			
先天性代謝異常による腎不全	146	3	149		28	177
(%)	(98.0)	(2.0)	(100.0)			
腎・尿路結核	126	2	128		30	158
(%)	(98.4)	(1.6)	(100.0)			
腎・尿路結石	349	5	354		76	430
(%)	(98.6)	(1.4)	(100.0)			
腎・尿路腫瘍	551	3	554	1	131	686
(%)	(99.5)	(0.5)	(100.0)			
閉塞性尿路障害	427	18	445		72	517
(%)	(96.0)	(4.0)	(100.0)			
骨髄腫	163	1	164		26	190
(%)	(99.4)	(0.6)	(100.0)			
腎形成不全	350	7	357		62	419
(%)	(98.0)	(2.0)	(100.0)			
不明	15,984	295	16,279	23	3,823	20,125
(%)	(98.2)	(1.8)	(100.0)			
再導入	1,155	31	1,186		231	1,417
(%)	(97.4)	(2.6)	(100.0)			
その他	4,439	111	4,550	8	956	5,514
(%)	(97.6)	(2.4)	(100.0)			
合計	182,127	7,089	189,216	260	39,495	228,971
(%)	(96.3)	(3.7)	(100.0)			
記載なし						
(%)						
総計	182,127	7,089	189,216	260	39,495	228,971
(%)	(96.3)	(3.7)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査