

表 1154 四肢切断の有無 身長(cm)別（施設血液透析患者 透析回数週3回）

身長	切断なし	切断あり	合計	不明	記載なし	総計
130未満	123	91	214	2	21	237
(%)	(57.5)	(42.5)	(100.0)			
130～	464	39	503		65	568
(%)	(92.2)	(7.8)	(100.0)			
135～	1,894	52	1,946	6	353	2,305
(%)	(97.3)	(2.7)	(100.0)			
140～	6,892	153	7,045	13	1,225	8,283
(%)	(97.8)	(2.2)	(100.0)			
145～	15,616	371	15,987	28	2,804	18,819
(%)	(97.7)	(2.3)	(100.0)			
150～	24,622	716	25,338	33	4,643	30,014
(%)	(97.2)	(2.8)	(100.0)			
155～	28,673	933	29,606	38	5,213	34,857
(%)	(96.8)	(3.2)	(100.0)			
160～	34,455	1,319	35,774	52	6,432	42,258
(%)	(96.3)	(3.7)	(100.0)			
165～	30,662	1,242	31,904	37	5,635	37,576
(%)	(96.1)	(3.9)	(100.0)			
170～	19,237	934	20,171	16	3,640	23,827
(%)	(95.4)	(4.6)	(100.0)			
175～	6,862	409	7,271	4	1,274	8,549
(%)	(94.4)	(5.6)	(100.0)			
180～	1,776	151	1,927	4	323	2,254
(%)	(92.2)	(7.8)	(100.0)			
185～	271	26	297		54	351
(%)	(91.2)	(8.8)	(100.0)			
190～	43	6	49		8	57
(%)	(87.8)	(12.2)	(100.0)			
合計	171,590	6,442	178,032	233	31,690	209,955
(%)	(96.4)	(3.6)	(100.0)			
記載なし	10,537	647	11,184	27	7,805	19,016
(%)	(94.2)	(5.8)	(100.0)			
総計	182,127	7,089	189,216	260	39,495	228,971
(%)	(96.3)	(3.7)	(100.0)			
平均	159.88	161.46	159.94	157.73	159.97	159.94
標準偏差	9.51	11.60	9.60	11.18	9.50	9.59

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査