

表 1158 四肢切断の有無 Body Mass Index別（施設血液透析患者 透析回数週3回）

Body Mass Index	切断なし	切断あり	合計	不明	記載なし	総計
14未満	1,030	67	1,097	1	213	1,311
(%)	(93.9)	(6.1)	(100.0)			
14～	6,165	351	6,516	7	1,083	7,606
(%)	(94.6)	(5.4)	(100.0)			
16～	20,038	816	20,854	18	3,608	24,480
(%)	(96.1)	(3.9)	(100.0)			
18～	35,666	1,155	36,821	39	6,296	43,156
(%)	(96.9)	(3.1)	(100.0)			
20～	39,319	1,210	40,529	67	7,084	47,680
(%)	(97.0)	(3.0)	(100.0)			
22～	30,199	1,044	31,243	43	5,524	36,810
(%)	(96.7)	(3.3)	(100.0)			
24～	18,021	641	18,662	27	3,224	21,913
(%)	(96.6)	(3.4)	(100.0)			
26～	9,484	438	9,922	17	1,749	11,688
(%)	(95.6)	(4.4)	(100.0)			
28～	4,886	238	5,124	3	864	5,991
(%)	(95.4)	(4.6)	(100.0)			
30～	4,186	263	4,449	7	708	5,164
(%)	(94.1)	(5.9)	(100.0)			
合計	168,994	6,223	175,217	229	30,353	205,799
(%)	(96.4)	(3.6)	(100.0)			
記載なし	13,133	866	13,999	31	9,142	23,172
(%)	(93.8)	(6.2)	(100.0)			
総計	182,127	7,089	189,216	260	39,495	228,971
(%)	(96.3)	(3.7)	(100.0)			
平均	21.46	21.62	21.46	21.81	21.46	21.46
標準偏差	3.64	4.18	3.66	3.43	3.63	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査