

表 1170 四肢切断の有無 PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	切断なし	切断あり	合計	不明	記載なし	総計
20未満	1,232	54	1,286		213	1,499
(%)	(95.8)	(4.2)	(100.0)			
20～	1,567	56	1,623		225	1,848
(%)	(96.5)	(3.5)	(100.0)			
35～	1,341	52	1,393		220	1,613
(%)	(96.3)	(3.7)	(100.0)			
50～	2,038	82	2,120		339	2,459
(%)	(96.1)	(3.9)	(100.0)			
75～	1,752	78	1,830		318	2,148
(%)	(95.7)	(4.3)	(100.0)			
100～	1,468	45	1,513		274	1,787
(%)	(97.0)	(3.0)	(100.0)			
125～	940	33	973		214	1,187
(%)	(96.6)	(3.4)	(100.0)			
150～	30	2	32		5	37
(%)	(93.8)	(6.3)	(100.0)			
151～	768	30	798		136	934
(%)	(96.2)	(3.8)	(100.0)			
180～	547	21	568		98	666
(%)	(96.3)	(3.7)	(100.0)			
210～	380	13	393		79	472
(%)	(96.7)	(3.3)	(100.0)			
240～	433	10	443		83	526
(%)	(97.7)	(2.3)	(100.0)			
300～	168	3	171		34	205
(%)	(98.2)	(1.8)	(100.0)			
350～	108	7	115		20	135
(%)	(93.9)	(6.1)	(100.0)			
400～	132	1	133		19	152
(%)	(99.2)	(0.8)	(100.0)			
500～	143	8	151		15	166
(%)	(94.7)	(5.3)	(100.0)			
合計	13,047	495	13,542		2,292	15,834
(%)	(96.3)	(3.7)	(100.0)			
記載なし	235	21	256		182	438
(%)	(91.8)	(8.2)	(100.0)			
総計	13,282	516	13,798		2,474	16,272
(%)	(96.3)	(3.7)	(100.0)			
平均	105.72	99.72	105.50		106.86	105.70
標準偏差	112.23	101.55	111.86		92.31	109.24

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査