

表 1171 四肢切断の有無 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン 濃度	切断なし	切断あり	合計	不明	記載なし	総計
5未満	220	11	231		30	261
(%)	(95.2)	(4.8)	(100.0)			
5～	302	12	314		48	362
(%)	(96.2)	(3.8)	(100.0)			
6～	745	38	783	3	156	942
(%)	(95.1)	(4.9)	(100.0)			
7～	2,855	155	3,010	6	562	3,578
(%)	(94.9)	(5.1)	(100.0)			
8～	9,451	494	9,945	22	1,759	11,726
(%)	(95.0)	(5.0)	(100.0)			
9～	29,887	1,270	31,157	46	5,835	37,038
(%)	(95.9)	(4.1)	(100.0)			
10～	61,076	2,143	63,219	78	11,920	75,217
(%)	(96.6)	(3.4)	(100.0)			
11～	49,945	1,730	51,675	63	9,741	61,479
(%)	(96.7)	(3.3)	(100.0)			
12～	18,079	722	18,801	27	3,542	22,370
(%)	(96.2)	(3.8)	(100.0)			
13～	4,440	217	4,657	4	876	5,537
(%)	(95.3)	(4.7)	(100.0)			
14～	1,564	85	1,649	1	283	1,933
(%)	(94.8)	(5.2)	(100.0)			
合計	178,564	6,877	185,441	250	34,752	220,443
(%)	(96.3)	(3.7)	(100.0)			
記載なし	3,563	212	3,775	10	4,743	8,528
(%)	(94.4)	(5.6)	(100.0)			
総計	182,127	7,089	189,216	260	39,495	228,971
(%)	(96.3)	(3.7)	(100.0)			
平均	10.69	10.63	10.69	10.51	10.70	10.69
標準偏差	1.28	1.40	1.28	1.29	1.26	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査