

表 1177 四肢切断の有無 透析前脈拍(拍/分)別 (施設血液透析患者 透析回数週3回)

透析前脈拍	切断なし	切断あり	合計	不明	記載なし	総計
40未満	115	15	130		25	155
(%)	(88.5)	(11.5)	(100.0)			
40～	1,585	54	1,639		290	1,929
(%)	(96.7)	(3.3)	(100.0)			
50～	13,086	363	13,449	16	2,308	15,773
(%)	(97.3)	(2.7)	(100.0)			
60～	40,091	1,217	41,308	38	7,395	48,741
(%)	(97.1)	(2.9)	(100.0)			
70～	47,544	1,732	49,276	62	8,846	58,184
(%)	(96.5)	(3.5)	(100.0)			
80～	29,441	1,357	30,798	57	5,868	36,723
(%)	(95.6)	(4.4)	(100.0)			
90～	12,723	701	13,424	16	2,478	15,918
(%)	(94.8)	(5.2)	(100.0)			
100～	3,974	212	4,186	6	712	4,904
(%)	(94.9)	(5.1)	(100.0)			
110～	939	50	989	2	166	1,157
(%)	(94.9)	(5.1)	(100.0)			
120～	421	17	438		60	498
(%)	(96.1)	(3.9)	(100.0)			
合計	149,919	5,718	155,637	197	28,148	183,982
(%)	(96.3)	(3.7)	(100.0)			
記載なし	32,208	1,371	33,579	63	11,347	44,989
(%)	(95.9)	(4.1)	(100.0)			
総計	182,127	7,089	189,216	260	39,495	228,971
(%)	(96.3)	(3.7)	(100.0)			
平均	74.37	76.97	74.47	76.62	74.65	74.50
標準偏差	12.80	13.26	12.83	12.18	12.62	12.80

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査