

表 1190 四肢切断の有無 身長(cm)別（血液透析濾過患者 透析回数週3回）

身長	切断なし	切断あり	合計	不明	記載なし	総計
130未満	26	9	35		3	38
(%)	(74.3)	(25.7)	(100.0)			
130～	47	5	52		18	70
(%)	(90.4)	(9.6)	(100.0)			
135～	243	2	245		44	289
(%)	(99.2)	(0.8)	(100.0)			
140～	955	17	972	1	190	1,163
(%)	(98.3)	(1.7)	(100.0)			
145～	2,428	59	2,487	1	507	2,995
(%)	(97.6)	(2.4)	(100.0)			
150～	4,311	128	4,439		773	5,212
(%)	(97.1)	(2.9)	(100.0)			
155～	5,048	137	5,185	3	891	6,079
(%)	(97.4)	(2.6)	(100.0)			
160～	5,908	181	6,089	2	1,086	7,177
(%)	(97.0)	(3.0)	(100.0)			
165～	5,506	241	5,747	1	974	6,722
(%)	(95.8)	(4.2)	(100.0)			
170～	3,889	187	4,076		709	4,785
(%)	(95.4)	(4.6)	(100.0)			
175～	1,458	61	1,519	1	266	1,786
(%)	(96.0)	(4.0)	(100.0)			
180～	398	26	424		69	493
(%)	(93.9)	(6.1)	(100.0)			
185～	63	3	66		11	77
(%)	(95.5)	(4.5)	(100.0)			
190～	8		8		1	9
(%)	(100.0)		(100.0)			
合計	30,288	1,056	31,344	9	5,542	36,895
(%)	(96.6)	(3.4)	(100.0)			
記載なし	1,471	88	1,559		1,492	3,051
(%)	(94.4)	(5.6)	(100.0)			
総計	31,759	1,144	32,903	9	7,034	39,946
(%)	(96.5)	(3.5)	(100.0)			
平均	160.73	162.56	160.79	158.11	160.46	160.74
標準偏差	9.48	10.24	9.51	11.07	9.68	9.54

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査