

表 1191 四肢切断の有無 透析後体重(kg)別（血液透析濾過患者 透析回数週3回）

透析後体重	切断なし	切断あり	合計	不明	記載なし	総計
20未満	8		8		4	12
(%)	(100.0)		(100.0)			
20～	61	2	63		13	76
(%)	(96.8)	(3.2)	(100.0)			
30～	2,158	71	2,229	1	452	2,682
(%)	(96.8)	(3.2)	(100.0)			
40～	8,112	235	8,347	3	1,655	10,005
(%)	(97.2)	(2.8)	(100.0)			
50～	10,061	351	10,412	1	2,077	12,490
(%)	(96.6)	(3.4)	(100.0)			
60～	6,736	248	6,984	2	1,379	8,365
(%)	(96.4)	(3.6)	(100.0)			
70～	2,771	120	2,891		532	3,423
(%)	(95.8)	(4.2)	(100.0)			
80～	1,035	56	1,091	1	181	1,273
(%)	(94.9)	(5.1)	(100.0)			
90～	349	31	380		75	455
(%)	(91.8)	(8.2)	(100.0)			
100～	107	9	116		26	142
(%)	(92.2)	(7.8)	(100.0)			
110～	51	1	52		9	61
(%)	(98.1)	(1.9)	(100.0)			
120～	31	1	32		5	37
(%)	(96.9)	(3.1)	(100.0)			
合計	31,480	1,125	32,605	8	6,408	39,021
(%)	(96.5)	(3.5)	(100.0)			
記載なし	279	19	298	1	626	925
(%)	(93.6)	(6.4)	(100.0)			
総計	31,759	1,144	32,903	9	7,034	39,946
(%)	(96.5)	(3.5)	(100.0)			
平均	56.61	59.05	56.70	55.64	56.28	56.63
標準偏差	13.01	14.34	13.06	15.10	12.87	13.03

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査