

表 1207 四肢切断の有無 透析前ヘモグロビン濃度(g/dL)別（血液透析濾過患者 透析回数週3回）

透析前ヘモグロビン 濃度	切断なし	切断あり	合計	不明	記載なし	総計
5未満	22	2	24		2	26
(%)	(91.7)	(8.3)	(100.0)			
5～	26	2	28		3	31
(%)	(92.9)	(7.1)	(100.0)			
6～	95	4	99		15	114
(%)	(96.0)	(4.0)	(100.0)			
7～	363	27	390		55	445
(%)	(93.1)	(6.9)	(100.0)			
8～	1,319	80	1,399	1	237	1,637
(%)	(94.3)	(5.7)	(100.0)			
9～	4,705	181	4,886		964	5,850
(%)	(96.3)	(3.7)	(100.0)			
10～	10,477	306	10,783	1	2,163	12,947
(%)	(97.2)	(2.8)	(100.0)			
11～	9,386	315	9,701	4	1,872	11,577
(%)	(96.8)	(3.2)	(100.0)			
12～	3,571	134	3,705	2	675	4,382
(%)	(96.4)	(3.6)	(100.0)			
13～	966	42	1,008		177	1,185
(%)	(95.8)	(4.2)	(100.0)			
14～	433	27	460		63	523
(%)	(94.1)	(5.9)	(100.0)			
合計	31,363	1,120	32,483	8	6,226	38,717
(%)	(96.6)	(3.4)	(100.0)			
記載なし	396	24	420	1	808	1,229
(%)	(94.3)	(5.7)	(100.0)			
総計	31,759	1,144	32,903	9	7,034	39,946
(%)	(96.5)	(3.5)	(100.0)			
平均	10.84	10.79	10.84	11.25	10.83	10.84
標準偏差	1.27	1.50	1.28	1.30	1.19	1.26

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査