

表 1247 大腿骨頭部骨折の既往 原疾患別（透析患者全体）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	70,668	2,799	73,467	240	23,263	96,970
(%)	(96.2)	(3.8)	(100.0)			
慢性腎盂腎炎	2,239	116	2,355	3	684	3,042
(%)	(95.1)	(4.9)	(100.0)			
急速進行性糸球体腎炎	1,824	72	1,896	7	591	2,494
(%)	(96.2)	(3.8)	(100.0)			
妊娠腎/妊娠中毒症	1,182	72	1,254	7	353	1,614
(%)	(94.3)	(5.7)	(100.0)			
その他の分類不能の腎炎	981	45	1,026	2	345	1,373
(%)	(95.6)	(4.4)	(100.0)			
多発性嚢胞腎	8,198	246	8,444	30	2,532	11,006
(%)	(97.1)	(2.9)	(100.0)			
腎硬化症	20,987	993	21,980	132	6,186	28,298
(%)	(95.5)	(4.5)	(100.0)			
悪性高血圧	1,875	61	1,936	7	630	2,573
(%)	(96.8)	(3.2)	(100.0)			
糖尿病性腎症	85,180	4,151	89,331	340	28,410	118,081
(%)	(95.4)	(4.6)	(100.0)			
SLE腎炎	1,616	118	1,734	8	494	2,236
(%)	(93.2)	(6.8)	(100.0)			
アミロイド腎	308	22	330	3	118	451
(%)	(93.3)	(6.7)	(100.0)			
痛風腎	860	22	882	6	225	1,113
(%)	(97.5)	(2.5)	(100.0)			
先天性代謝異常による腎不全	199	4	203	1	68	272
(%)	(98.0)	(2.0)	(100.0)			
腎・尿路結核	142	11	153		53	206
(%)	(92.8)	(7.2)	(100.0)			
腎・尿路結石	421	18	439	1	141	581
(%)	(95.9)	(4.1)	(100.0)			
腎・尿路腫瘍	646	13	659	1	226	886
(%)	(98.0)	(2.0)	(100.0)			
閉塞性尿路障害	547	21	568		155	723
(%)	(96.3)	(3.7)	(100.0)			
骨髄腫	204	8	212	1	63	276
(%)	(96.2)	(3.8)	(100.0)			
腎形成不全	500	8	508	1	140	649
(%)	(98.4)	(1.6)	(100.0)			
不明	19,182	1,007	20,189	130	7,225	27,544
(%)	(95.0)	(5.0)	(100.0)			
再導入	1,599	61	1,660	2	501	2,163
(%)	(96.3)	(3.7)	(100.0)			
その他	5,376	255	5,631	34	1,892	7,557
(%)	(95.5)	(4.5)	(100.0)			
合計	224,734	10,123	234,857	956	74,295	310,108
(%)	(95.7)	(4.3)	(100.0)			
記載なし						
(%)						
総計	224,734	10,123	234,857	956	74,295	310,108
(%)	(95.7)	(4.3)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査