

表 1255 大腿骨頸部骨折の既往 身長(cm)別（透析患者全体）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	285	25	310	2	42	354
(%)	(91.9)	(8.1)	(100.0)			
130～	521	75	596	2	98	696
(%)	(87.4)	(12.6)	(100.0)			
135～	2,122	258	2,380	18	439	2,837
(%)	(89.2)	(10.8)	(100.0)			
140～	7,765	805	8,570	47	1,680	10,297
(%)	(90.6)	(9.4)	(100.0)			
145～	18,088	1,484	19,572	72	4,029	23,673
(%)	(92.4)	(7.6)	(100.0)			
150～	29,570	1,876	31,446	126	6,663	38,235
(%)	(94.0)	(6.0)	(100.0)			
155～	35,129	1,597	36,726	142	7,604	44,472
(%)	(95.7)	(4.3)	(100.0)			
160～	42,689	1,420	44,109	155	9,386	53,650
(%)	(96.8)	(3.2)	(100.0)			
165～	38,546	1,003	39,549	150	8,289	47,988
(%)	(97.5)	(2.5)	(100.0)			
170～	24,851	603	25,454	96	5,524	31,074
(%)	(97.6)	(2.4)	(100.0)			
175～	9,123	205	9,328	26	1,965	11,319
(%)	(97.8)	(2.2)	(100.0)			
180～	2,421	45	2,466	11	502	2,979
(%)	(98.2)	(1.8)	(100.0)			
185～	371	12	383	3	79	465
(%)	(96.9)	(3.1)	(100.0)			
190～	59	3	62		15	77
(%)	(95.2)	(4.8)	(100.0)			
合計	211,540	9,411	220,951	850	46,315	268,116
(%)	(95.7)	(4.3)	(100.0)			
記載なし	13,194	712	13,906	106	27,980	41,992
(%)	(94.9)	(5.1)	(100.0)			
総計	224,734	10,123	234,857	956	74,295	310,108
(%)	(95.7)	(4.3)	(100.0)			
平均	160.22	155.44	160.02	159.32	160.18	160.05
標準偏差	9.61	9.95	9.67	10.43	9.54	9.65

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査