

表 1257 大腿骨頸部骨折の既往 身長(cm)別 (透析患者全体 女性のみ)

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	178	21	199	1	27	227
(%)	(89.4)	(10.6)	(100.0)			
130～	474	74	548	2	87	637
(%)	(86.5)	(13.5)	(100.0)			
135～	2,028	252	2,280	18	429	2,727
(%)	(88.9)	(11.1)	(100.0)			
140～	7,393	775	8,168	44	1,604	9,816
(%)	(90.5)	(9.5)	(100.0)			
145～	16,548	1,403	17,951	65	3,696	21,712
(%)	(92.2)	(7.8)	(100.0)			
150～	22,541	1,609	24,150	92	5,144	29,386
(%)	(93.3)	(6.7)	(100.0)			
155～	16,173	930	17,103	63	3,572	20,738
(%)	(94.6)	(5.4)	(100.0)			
160～	6,874	309	7,183	21	1,562	8,766
(%)	(95.7)	(4.3)	(100.0)			
165～	1,685	46	1,731	4	374	2,109
(%)	(97.3)	(2.7)	(100.0)			
170～	328	5	333		74	407
(%)	(98.5)	(1.5)	(100.0)			
175～	77	4	81		18	99
(%)	(95.1)	(4.9)	(100.0)			
180～	11	2	13		2	15
(%)	(84.6)	(15.4)	(100.0)			
185～	2	1	3			3
(%)	(66.7)	(33.3)	(100.0)			
190～					1	1
(%)						
合計	74,312	5,431	79,743	310	16,590	96,643
(%)	(93.2)	(6.8)	(100.0)			
記載なし	4,649	420	5,069	42	10,213	15,324
(%)	(91.7)	(8.3)	(100.0)			
総計	78,961	5,851	84,812	352	26,803	111,967
(%)	(93.1)	(6.9)	(100.0)			
平均	151.59	149.54	151.45	150.25	151.67	151.48
標準偏差	7.01	6.92	7.02	6.96	6.89	7.00

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査