

表 1258 大腿骨頸部骨折の既往 Body Mass Index別（透析患者全体）

Body Mass Index	既往なし	既往あり	合計	不明	記載なし	総計
14未満	1,155	171	1,326	1	259	1,586
(%)	(87.1)	(12.9)	(100.0)			
14～	6,935	807	7,742	33	1,415	9,190
(%)	(89.6)	(10.4)	(100.0)			
16～	23,637	1,765	25,402	89	4,732	30,223
(%)	(93.1)	(6.9)	(100.0)			
18～	42,999	2,288	45,287	162	8,408	53,857
(%)	(94.9)	(5.1)	(100.0)			
20～	48,311	1,830	50,141	201	9,578	59,920
(%)	(96.4)	(3.6)	(100.0)			
22～	37,611	1,213	38,824	142	7,455	46,421
(%)	(96.9)	(3.1)	(100.0)			
24～	22,866	576	23,442	103	4,421	27,966
(%)	(97.5)	(2.5)	(100.0)			
26～	12,275	260	12,535	57	2,337	14,929
(%)	(97.9)	(2.1)	(100.0)			
28～	6,355	130	6,485	18	1,213	7,716
(%)	(98.0)	(2.0)	(100.0)			
30～	5,493	114	5,607	22	982	6,611
(%)	(98.0)	(2.0)	(100.0)			
合計	207,637	9,154	216,791	828	40,800	258,419
(%)	(95.8)	(4.2)	(100.0)			
記載なし	17,097	969	18,066	128	33,495	51,689
(%)	(94.6)	(5.4)	(100.0)			
総計	224,734	10,123	234,857	956	74,295	310,108
(%)	(95.7)	(4.3)	(100.0)			
平均	21.57	19.98	21.51	21.65	21.51	21.51
標準偏差	3.66	3.47	3.66	3.61	3.63	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査