

表 1267 大腿骨頸部骨折の既往 PTH値(pg/mL)別（透析患者全体）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	1,504	84	1,588	1	305	1,894
(%)	(94.7)	(5.3)	(100.0)			
20～	1,891	93	1,984		347	2,331
(%)	(95.3)	(4.7)	(100.0)			
35～	1,660	87	1,747		296	2,043
(%)	(95.0)	(5.0)	(100.0)			
50～	2,538	112	2,650		503	3,153
(%)	(95.8)	(4.2)	(100.0)			
75～	2,193	98	2,291		457	2,748
(%)	(95.7)	(4.3)	(100.0)			
100～	1,834	52	1,886	3	409	2,298
(%)	(97.2)	(2.8)	(100.0)			
125～	1,190	38	1,228	1	310	1,539
(%)	(96.9)	(3.1)	(100.0)			
150～	36	2	38		10	48
(%)	(94.7)	(5.3)	(100.0)			
151～	998	44	1,042		221	1,263
(%)	(95.8)	(4.2)	(100.0)			
180～	699	29	728		147	875
(%)	(96.0)	(4.0)	(100.0)			
210～	498	17	515		113	628
(%)	(96.7)	(3.3)	(100.0)			
240～	534	26	560		114	674
(%)	(95.4)	(4.6)	(100.0)			
300～	228	4	232		52	284
(%)	(98.3)	(1.7)	(100.0)			
350～	151	2	153		26	179
(%)	(98.7)	(1.3)	(100.0)			
400～	165	5	170		32	202
(%)	(97.1)	(2.9)	(100.0)			
500～	193	8	201		46	247
(%)	(96.0)	(4.0)	(100.0)			
合計	16,312	701	17,013	5	3,388	20,406
(%)	(95.9)	(4.1)	(100.0)			
記載なし	357	36	393		264	657
(%)	(90.8)	(9.2)	(100.0)			
総計	16,669	737	17,406	5	3,652	21,063
(%)	(95.8)	(4.2)	(100.0)			
平均	107.32	98.30	106.94	89.80	113.36	108.01
標準偏差	110.95	123.19	111.49	46.64	125.89	114.01

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査