

表 1268 大腿骨頸部骨折の既往 透析前ヘモグロビン濃度(g/dL)別（透析患者全体）

透析前ヘモグロビン濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	253	15	268		35	303
(%)	(94.4)	(5.6)	(100.0)			
5～	340	23	363	1	65	429
(%)	(93.7)	(6.3)	(100.0)			
6～	887	63	950	10	200	1,160
(%)	(93.4)	(6.6)	(100.0)			
7～	3,434	235	3,669	18	748	4,435
(%)	(93.6)	(6.4)	(100.0)			
8～	11,355	754	12,109	69	2,376	14,554
(%)	(93.8)	(6.2)	(100.0)			
9～	36,020	1,893	37,913	170	7,906	45,989
(%)	(95.0)	(5.0)	(100.0)			
10～	74,192	3,183	77,375	317	16,220	93,912
(%)	(95.9)	(4.1)	(100.0)			
11～	62,005	2,403	64,408	226	13,539	78,173
(%)	(96.3)	(3.7)	(100.0)			
12～	22,965	877	23,842	92	4,920	28,854
(%)	(96.3)	(3.7)	(100.0)			
13～	5,822	220	6,042	9	1,257	7,308
(%)	(96.4)	(3.6)	(100.0)			
14～	2,174	76	2,250	5	418	2,673
(%)	(96.6)	(3.4)	(100.0)			
合計	219,447	9,742	229,189	917	47,684	277,790
(%)	(95.7)	(4.3)	(100.0)			
記載なし	5,287	381	5,668	39	26,611	32,318
(%)	(93.3)	(6.7)	(100.0)			
総計	224,734	10,123	234,857	956	74,295	310,108
(%)	(95.7)	(4.3)	(100.0)			
平均	10.72	10.51	10.71	10.50	10.72	10.71
標準偏差	1.29	1.35	1.29	1.26	1.26	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査