

表 1274 大腿骨頸部骨折の既往 透析前脈拍(拍/分)別 (透析患者全体)

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	140	11	151		34	185
(%)	(92.7)	(7.3)	(100.0)			
40～	1,892	107	1,999		382	2,381
(%)	(94.6)	(5.4)	(100.0)			
50～	15,609	706	16,315	40	3,157	19,512
(%)	(95.7)	(4.3)	(100.0)			
60～	48,327	2,302	50,629	118	10,034	60,781
(%)	(95.5)	(4.5)	(100.0)			
70～	58,299	2,664	60,963	152	12,157	73,272
(%)	(95.6)	(4.4)	(100.0)			
80～	37,030	1,557	38,587	106	8,121	46,814
(%)	(96.0)	(4.0)	(100.0)			
90～	16,402	649	17,051	51	3,488	20,590
(%)	(96.2)	(3.8)	(100.0)			
100～	5,187	184	5,371	17	1,005	6,393
(%)	(96.6)	(3.4)	(100.0)			
110～	1,241	39	1,280	3	245	1,528
(%)	(97.0)	(3.0)	(100.0)			
120～	545	22	567	1	87	655
(%)	(96.1)	(3.9)	(100.0)			
合計	184,672	8,241	192,913	488	38,710	232,111
(%)	(95.7)	(4.3)	(100.0)			
記載なし	40,062	1,882	41,944	468	35,585	77,997
(%)	(95.5)	(4.5)	(100.0)			
総計	224,734	10,123	234,857	956	74,295	310,108
(%)	(95.7)	(4.3)	(100.0)			
平均	74.70	73.83	74.66	75.99	74.81	74.69
標準偏差	12.89	12.52	12.87	12.74	12.66	12.84

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査