

表 1286 大腿骨頸部骨折の既往 身長(cm)別（施設血液透析患者 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	186	20	206	2	29	237
(%)	(90.3)	(9.7)	(100.0)			
130～	433	68	501	1	66	568
(%)	(86.4)	(13.6)	(100.0)			
135～	1,733	213	1,946	16	343	2,305
(%)	(89.1)	(10.9)	(100.0)			
140～	6,359	671	7,030	40	1,213	8,283
(%)	(90.5)	(9.5)	(100.0)			
145～	14,707	1,215	15,922	63	2,834	18,819
(%)	(92.4)	(7.6)	(100.0)			
150～	23,616	1,528	25,144	111	4,759	30,014
(%)	(93.9)	(6.1)	(100.0)			
155～	28,039	1,295	29,334	123	5,400	34,857
(%)	(95.6)	(4.4)	(100.0)			
160～	34,270	1,152	35,422	133	6,703	42,258
(%)	(96.7)	(3.3)	(100.0)			
165～	30,707	820	31,527	128	5,921	37,576
(%)	(97.4)	(2.6)	(100.0)			
170～	19,387	491	19,878	80	3,869	23,827
(%)	(97.5)	(2.5)	(100.0)			
175～	7,006	167	7,173	24	1,352	8,549
(%)	(97.7)	(2.3)	(100.0)			
180～	1,868	31	1,899	10	345	2,254
(%)	(98.4)	(1.6)	(100.0)			
185～	287	8	295	1	55	351
(%)	(97.3)	(2.7)	(100.0)			
190～	46	3	49		8	57
(%)	(93.9)	(6.1)	(100.0)			
合計	168,644	7,682	176,326	732	32,897	209,955
(%)	(95.6)	(4.4)	(100.0)			
記載なし	10,383	551	10,934	71	8,011	19,016
(%)	(95.0)	(5.0)	(100.0)			
総計	179,027	8,233	187,260	803	40,908	228,971
(%)	(95.6)	(4.4)	(100.0)			
平均	160.12	155.35	159.91	159.20	160.12	159.94
標準偏差	9.53	9.96	9.60	10.44	9.49	9.59

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査