

表 1287 大腿骨頸部骨折の既往 身長(cm)別（施設血液透析患者 男性のみ 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	72	4	76	1	10	87
(%)	(94.7)	(5.3)	(100.0)			
130～	37	1	38		7	45
(%)	(97.4)	(2.6)	(100.0)			
135～	80	5	85		9	94
(%)	(94.1)	(5.9)	(100.0)			
140～	310	25	335	3	59	397
(%)	(92.5)	(7.5)	(100.0)			
145～	1,300	72	1,372	7	255	1,634
(%)	(94.8)	(5.2)	(100.0)			
150～	5,815	217	6,032	29	1,152	7,213
(%)	(96.4)	(3.6)	(100.0)			
155～	15,590	545	16,135	69	2,958	19,162
(%)	(96.6)	(3.4)	(100.0)			
160～	29,076	916	29,992	115	5,637	35,744
(%)	(96.9)	(3.1)	(100.0)			
165～	29,399	782	30,181	124	5,666	35,971
(%)	(97.4)	(2.6)	(100.0)			
170～	19,135	486	19,621	80	3,819	23,520
(%)	(97.5)	(2.5)	(100.0)			
175～	6,948	163	7,111	24	1,340	8,475
(%)	(97.7)	(2.3)	(100.0)			
180～	1,862	29	1,891	10	345	2,246
(%)	(98.5)	(1.5)	(100.0)			
185～	285	7	292	1	55	348
(%)	(97.6)	(2.4)	(100.0)			
190～	46	3	49		7	56
(%)	(93.9)	(6.1)	(100.0)			
合計	109,955	3,255	113,210	463	21,319	134,992
(%)	(97.1)	(2.9)	(100.0)			
記載なし	6,742	219	6,961	42	5,038	12,041
(%)	(96.9)	(3.1)	(100.0)			
総計	116,697	3,474	120,171	505	26,357	147,033
(%)	(97.1)	(2.9)	(100.0)			
平均	164.75	163.38	164.71	164.34	164.77	164.72
標準偏差	7.25	7.54	7.27	8.48	7.22	7.26

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査