

表 1288 大腿骨頸部骨折の既往 身長(cm)別（施設血液透析患者 女性のみ 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	114	16	130	1	19	150
(%)	(87.7)	(12.3)	(100.0)			
130～	396	67	463	1	59	523
(%)	(85.5)	(14.5)	(100.0)			
135～	1,653	208	1,861	16	334	2,211
(%)	(88.8)	(11.2)	(100.0)			
140～	6,049	646	6,695	37	1,154	7,886
(%)	(90.4)	(9.6)	(100.0)			
145～	13,407	1,143	14,550	56	2,579	17,185
(%)	(92.1)	(7.9)	(100.0)			
150～	17,801	1,311	19,112	82	3,607	22,801
(%)	(93.1)	(6.9)	(100.0)			
155～	12,449	750	13,199	54	2,442	15,695
(%)	(94.3)	(5.7)	(100.0)			
160～	5,194	236	5,430	18	1,066	6,514
(%)	(95.7)	(4.3)	(100.0)			
165～	1,308	38	1,346	4	255	1,605
(%)	(97.2)	(2.8)	(100.0)			
170～	252	5	257		50	307
(%)	(98.1)	(1.9)	(100.0)			
175～	58	4	62		12	74
(%)	(93.5)	(6.5)	(100.0)			
180～	6	2	8			8
(%)	(75.0)	(25.0)	(100.0)			
185～	2	1	3			3
(%)	(66.7)	(33.3)	(100.0)			
190～					1	1
(%)						
合計	58,689	4,427	63,116	269	11,578	74,963
(%)	(93.0)	(7.0)	(100.0)			
記載なし	3,641	332	3,973	29	2,973	6,975
(%)	(91.6)	(8.4)	(100.0)			
総計	62,330	4,759	67,089	298	14,551	81,938
(%)	(92.9)	(7.1)	(100.0)			
平均	151.45	149.45	151.31	150.35	151.55	151.34
標準偏差	6.87	6.95	6.89	6.98	6.82	6.89

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査