

表 1298 大腿骨頸部骨折の既往 Body Mass Index別（施設血液透析患者 透析回数週3回）

Body Mass Index	既往なし	既往あり	合計	不明	記載なし	総計
14未満	952	148	1,100	1	210	1,311
(%)	(86.5)	(13.5)	(100.0)			
14～	5,797	684	6,481	28	1,097	7,606
(%)	(89.4)	(10.6)	(100.0)			
16～	19,251	1,464	20,715	74	3,691	24,480
(%)	(92.9)	(7.1)	(100.0)			
18～	34,622	1,905	36,527	140	6,489	43,156
(%)	(94.8)	(5.2)	(100.0)			
20～	38,632	1,514	40,146	175	7,359	47,680
(%)	(96.2)	(3.8)	(100.0)			
22～	29,997	961	30,958	129	5,723	36,810
(%)	(96.9)	(3.1)	(100.0)			
24～	17,977	471	18,448	86	3,379	21,913
(%)	(97.4)	(2.6)	(100.0)			
26～	9,592	215	9,807	48	1,833	11,688
(%)	(97.8)	(2.2)	(100.0)			
28～	4,943	110	5,053	17	921	5,991
(%)	(97.8)	(2.2)	(100.0)			
30～	4,302	89	4,391	19	754	5,164
(%)	(98.0)	(2.0)	(100.0)			
合計	166,065	7,561	173,626	717	31,456	205,799
(%)	(95.6)	(4.4)	(100.0)			
記載なし	12,962	672	13,634	86	9,452	23,172
(%)	(95.1)	(4.9)	(100.0)			
総計	179,027	8,233	187,260	803	40,908	228,971
(%)	(95.6)	(4.4)	(100.0)			
平均	21.52	19.93	21.46	21.66	21.49	21.46
標準偏差	3.65	3.47	3.66	3.61	3.63	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査