

表 1301 大腿骨頸部骨折の既往 透析前BUN(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前BUN	既往なし	既往あり	合計	不明	記載なし	総計
20未満	349	45	394	2	83	479
(%)	(88.6)	(11.4)	(100.0)			
20～	1,827	222	2,049	13	418	2,480
(%)	(89.2)	(10.8)	(100.0)			
30～	7,862	696	8,558	42	1,758	10,358
(%)	(91.9)	(8.1)	(100.0)			
40～	20,910	1,387	22,297	81	4,433	26,811
(%)	(93.8)	(6.2)	(100.0)			
50～	33,501	1,762	35,263	124	7,272	42,659
(%)	(95.0)	(5.0)	(100.0)			
60～	33,806	1,437	35,243	135	7,265	42,643
(%)	(95.9)	(4.1)	(100.0)			
70～	22,652	838	23,490	96	4,736	28,322
(%)	(96.4)	(3.6)	(100.0)			
80～	10,570	370	10,940	53	2,219	13,212
(%)	(96.6)	(3.4)	(100.0)			
90～	3,833	129	3,962	15	826	4,803
(%)	(96.7)	(3.3)	(100.0)			
100～	1,095	44	1,139	6	203	1,348
(%)	(96.1)	(3.9)	(100.0)			
110～	315	13	328	1	65	394
(%)	(96.0)	(4.0)	(100.0)			
120～	161	5	166	3	35	204
(%)	(97.0)	(3.0)	(100.0)			
合計	136,881	6,948	143,829	571	29,313	173,713
(%)	(95.2)	(4.8)	(100.0)			
記載なし	1,680	115	1,795	14	2,687	4,496
(%)	(93.6)	(6.4)	(100.0)			
総計	138,561	7,063	145,624	585	32,000	178,209
(%)	(95.1)	(4.9)	(100.0)			
平均	61.17	56.53	60.94	61.36	60.92	60.94
標準偏差	15.59	16.14	15.65	17.01	15.57	15.64

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査