

表 1317 大腿骨頸部骨折の既往 透析前補正カルシウム濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

透析前補正 カルシウム濃度	既往なし	既往あり	合計	不明	記載なし	総計
7.0未満	711	24	735	3	136	874
(%)	(96.7)	(3.3)	(100.0)			
7.0～	1,000	46	1,046	7	213	1,266
(%)	(95.6)	(4.4)	(100.0)			
7.5～	3,889	166	4,055	21	895	4,971
(%)	(95.9)	(4.1)	(100.0)			
8.0～	12,917	530	13,447	51	2,792	16,290
(%)	(96.1)	(3.9)	(100.0)			
8.4～	3,768	151	3,919	17	740	4,676
(%)	(96.1)	(3.9)	(100.0)			
8.5～	42,894	1,778	44,672	190	9,040	53,902
(%)	(96.0)	(4.0)	(100.0)			
9.0～	52,136	2,353	54,489	248	10,499	65,236
(%)	(95.7)	(4.3)	(100.0)			
9.5～	33,801	1,635	35,436	128	6,840	42,404
(%)	(95.4)	(4.6)	(100.0)			
10.0～	4,198	193	4,391	21	815	5,227
(%)	(95.6)	(4.4)	(100.0)			
10.1～	10,444	558	11,002	49	2,226	13,277
(%)	(94.9)	(5.1)	(100.0)			
10.5～	5,067	301	5,368	24	1,061	6,453
(%)	(94.4)	(5.6)	(100.0)			
11.0～	1,676	140	1,816	7	342	2,165
(%)	(92.3)	(7.7)	(100.0)			
11.5～	592	40	632	3	134	769
(%)	(93.7)	(6.3)	(100.0)			
12.0～	428	23	451	4	84	539
(%)	(94.9)	(5.1)	(100.0)			
合計	173,521	7,938	181,459	773	35,817	218,049
(%)	(95.6)	(4.4)	(100.0)			
記載なし	5,506	295	5,801	30	5,091	10,922
(%)	(94.9)	(5.1)	(100.0)			
総計	179,027	8,233	187,260	803	40,908	228,971
(%)	(95.6)	(4.4)	(100.0)			
平均	9.18	9.26	9.19	9.19	9.17	9.18
標準偏差	0.74	0.77	0.74	0.78	0.74	0.74

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査