

表 1320 大腿骨頸部骨折の既往 PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	1,202	66	1,268	1	230	1,499
(%)	(94.8)	(5.2)	(100.0)			
20～	1,536	79	1,615		233	1,848
(%)	(95.1)	(4.9)	(100.0)			
35～	1,316	73	1,389		224	1,613
(%)	(94.7)	(5.3)	(100.0)			
50～	2,018	92	2,110		349	2,459
(%)	(95.6)	(4.4)	(100.0)			
75～	1,745	81	1,826		322	2,148
(%)	(95.6)	(4.4)	(100.0)			
100～	1,453	41	1,494	3	290	1,787
(%)	(97.3)	(2.7)	(100.0)			
125～	926	29	955	1	231	1,187
(%)	(97.0)	(3.0)	(100.0)			
150～	28	2	30		7	37
(%)	(93.3)	(6.7)	(100.0)			
151～	750	36	786		148	934
(%)	(95.4)	(4.6)	(100.0)			
180～	537	25	562		104	666
(%)	(95.6)	(4.4)	(100.0)			
210～	373	9	382		90	472
(%)	(97.6)	(2.4)	(100.0)			
240～	423	19	442		84	526
(%)	(95.7)	(4.3)	(100.0)			
300～	166	3	169		36	205
(%)	(98.2)	(1.8)	(100.0)			
350～	112	2	114		21	135
(%)	(98.2)	(1.8)	(100.0)			
400～	129	4	133		19	152
(%)	(97.0)	(3.0)	(100.0)			
500～	135	7	142		24	166
(%)	(95.1)	(4.9)	(100.0)			
合計	12,849	568	13,417	5	2,412	15,834
(%)	(95.8)	(4.2)	(100.0)			
記載なし	224	31	255		183	438
(%)	(87.8)	(12.2)	(100.0)			
総計	13,073	599	13,672	5	2,595	16,272
(%)	(95.6)	(4.4)	(100.0)			
平均	105.27	97.72	104.95	89.80	109.86	105.70
標準偏差	109.27	129.84	110.22	46.64	103.65	109.24

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査