

表 1321 大腿骨頸部骨折の既往 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	217	13	230		31	261
(%)	(94.3)	(5.7)	(100.0)			
5～	292	19	311	1	50	362
(%)	(93.9)	(6.1)	(100.0)			
6～	720	52	772	9	161	942
(%)	(93.3)	(6.7)	(100.0)			
7～	2,793	196	2,989	15	574	3,578
(%)	(93.4)	(6.6)	(100.0)			
8～	9,247	622	9,869	53	1,804	11,726
(%)	(93.7)	(6.3)	(100.0)			
9～	29,266	1,593	30,859	145	6,034	37,038
(%)	(94.8)	(5.2)	(100.0)			
10～	60,012	2,642	62,654	275	12,288	75,217
(%)	(95.8)	(4.2)	(100.0)			
11～	49,226	1,954	51,180	198	10,101	61,479
(%)	(96.2)	(3.8)	(100.0)			
12～	17,905	730	18,635	75	3,660	22,370
(%)	(96.1)	(3.9)	(100.0)			
13～	4,436	174	4,610	6	921	5,537
(%)	(96.2)	(3.8)	(100.0)			
14～	1,578	59	1,637	3	293	1,933
(%)	(96.4)	(3.6)	(100.0)			
合計	175,692	8,054	183,746	780	35,917	220,443
(%)	(95.6)	(4.4)	(100.0)			
記載なし	3,335	179	3,514	23	4,991	8,528
(%)	(94.9)	(5.1)	(100.0)			
総計	179,027	8,233	187,260	803	40,908	228,971
(%)	(95.6)	(4.4)	(100.0)			
平均	10.70	10.50	10.69	10.50	10.70	10.69
標準偏差	1.28	1.34	1.28	1.23	1.26	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査