

表 1339 大腿骨頸部骨折の既往 身長(cm)別（血液透析濾過患者 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	30	4	34		4	38
(%)	(88.2)	(11.8)	(100.0)			
130～	46	4	50		20	70
(%)	(92.0)	(8.0)	(100.0)			
135～	227	20	247		42	289
(%)	(91.9)	(8.1)	(100.0)			
140～	887	84	971	3	189	1,163
(%)	(91.3)	(8.7)	(100.0)			
145～	2,295	167	2,462	4	529	2,995
(%)	(93.2)	(6.8)	(100.0)			
150～	4,141	234	4,375	4	833	5,212
(%)	(94.7)	(5.3)	(100.0)			
155～	4,914	194	5,108	7	964	6,079
(%)	(96.2)	(3.8)	(100.0)			
160～	5,854	172	6,026	5	1,146	7,177
(%)	(97.1)	(2.9)	(100.0)			
165～	5,545	122	5,667	5	1,050	6,722
(%)	(97.8)	(2.2)	(100.0)			
170～	3,967	65	4,032	4	749	4,785
(%)	(98.4)	(1.6)	(100.0)			
175～	1,471	27	1,498	1	287	1,786
(%)	(98.2)	(1.8)	(100.0)			
180～	412	9	421		72	493
(%)	(97.9)	(2.1)	(100.0)			
185～	61	3	64	1	12	77
(%)	(95.3)	(4.7)	(100.0)			
190～	7		7		2	9
(%)	(100.0)		(100.0)			
合計	29,857	1,105	30,962	34	5,899	36,895
(%)	(96.4)	(3.6)	(100.0)			
記載なし	1,321	55	1,376	5	1,670	3,051
(%)	(96.0)	(4.0)	(100.0)			
総計	31,178	1,160	32,338	39	7,569	39,946
(%)	(96.4)	(3.6)	(100.0)			
平均	160.95	156.04	160.78	159.32	160.53	160.74
標準偏差	9.46	9.69	9.51	10.74	9.65	9.54

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査