

表 1370 大腿骨頸部骨折の既往 透析前補正カルシウム濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前補正 カルシウム濃度	既往なし	既往あり	合計	不明	記載なし	総計
7.0未満	135	4	139		21	160
(%)	(97.1)	(2.9)	(100.0)			
7.0～	213	10	223	1	35	259
(%)	(95.5)	(4.5)	(100.0)			
7.5～	744	35	779	1	171	951
(%)	(95.5)	(4.5)	(100.0)			
8.0～	2,307	79	2,386		482	2,868
(%)	(96.7)	(3.3)	(100.0)			
8.4～	676	24	700	2	116	818
(%)	(96.6)	(3.4)	(100.0)			
8.5～	7,481	253	7,734	8	1,559	9,301
(%)	(96.7)	(3.3)	(100.0)			
9.0～	9,234	307	9,541	15	1,959	11,515
(%)	(96.8)	(3.2)	(100.0)			
9.5～	6,011	231	6,242	4	1,366	7,612
(%)	(96.3)	(3.7)	(100.0)			
10.0～	696	29	725		178	903
(%)	(96.0)	(4.0)	(100.0)			
10.1～	1,941	85	2,026	2	424	2,452
(%)	(95.8)	(4.2)	(100.0)			
10.5～	884	49	933		237	1,170
(%)	(94.7)	(5.3)	(100.0)			
11.0～	298	15	313	1	69	383
(%)	(95.2)	(4.8)	(100.0)			
11.5～	101	9	110		22	132
(%)	(91.8)	(8.2)	(100.0)			
12.0～	56		56	1	14	71
(%)	(100.0)		(100.0)			
合計	30,777	1,130	31,907	35	6,653	38,595
(%)	(96.5)	(3.5)	(100.0)			
記載なし	401	30	431	4	916	1,351
(%)	(93.0)	(7.0)	(100.0)			
総計	31,178	1,160	32,338	39	7,569	39,946
(%)	(96.4)	(3.6)	(100.0)			
平均	9.18	9.24	9.18	9.26	9.22	9.19
標準偏差	0.74	0.79	0.75	0.92	0.75	0.75

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査