

表 1380 大腿骨頸部骨折の既往 透析前脈拍(拍/分)別 (血液透析濾過患者 透析回数週3回)

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	13	1	14		3	17
(%)	(92.9)	(7.1)	(100.0)			
40～	241	14	255		40	295
(%)	(94.5)	(5.5)	(100.0)			
50～	2,094	74	2,168	1	463	2,632
(%)	(96.6)	(3.4)	(100.0)			
60～	6,729	262	6,991	6	1,493	8,490
(%)	(96.3)	(3.7)	(100.0)			
70～	8,377	314	8,691	9	1,845	10,545
(%)	(96.4)	(3.6)	(100.0)			
80～	5,577	202	5,779	6	1,278	7,063
(%)	(96.5)	(3.5)	(100.0)			
90～	2,599	82	2,681	2	610	3,293
(%)	(96.9)	(3.1)	(100.0)			
100～	829	29	858	1	183	1,042
(%)	(96.6)	(3.4)	(100.0)			
110～	218	4	222		39	261
(%)	(98.2)	(1.8)	(100.0)			
120～	93	2	95		11	106
(%)	(97.9)	(2.1)	(100.0)			
合計	26,770	984	27,754	25	5,965	33,744
(%)	(96.5)	(3.5)	(100.0)			
記載なし	4,408	176	4,584	14	1,604	6,202
(%)	(96.2)	(3.8)	(100.0)			
総計	31,178	1,160	32,338	39	7,569	39,946
(%)	(96.4)	(3.6)	(100.0)			
平均	75.44	74.60	75.41	77.20	75.52	75.43
標準偏差	13.00	12.60	12.99	11.49	12.67	12.93

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査