

表 1405 大腿骨頸部骨折の既往 PTH値(pg/mL)別（腹膜透析患者）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	37	3	40			40
(%)	(92.5)	(7.5)	(100.0)			
20～	45		45			45
(%)	(100.0)		(100.0)			
35～	53	1	54		1	55
(%)	(98.1)	(1.9)	(100.0)			
50～	75	1	76		1	77
(%)	(98.7)	(1.3)	(100.0)			
75～	68	1	69			69
(%)	(98.6)	(1.4)	(100.0)			
100～	70	1	71			71
(%)	(98.6)	(1.4)	(100.0)			
125～	45		45			45
(%)	(100.0)		(100.0)			
150～	1		1			1
(%)	(100.0)		(100.0)			
151～	42		42		1	43
(%)	(100.0)		(100.0)			
180～	32		32		2	34
(%)	(100.0)		(100.0)			
210～	26	1	27			27
(%)	(96.3)	(3.7)	(100.0)			
240～	22	1	23			23
(%)	(95.7)	(4.3)	(100.0)			
300～	19		19			19
(%)	(100.0)		(100.0)			
350～	8		8			8
(%)	(100.0)		(100.0)			
400～	11		11			11
(%)	(100.0)		(100.0)			
500～	11		11			11
(%)	(100.0)		(100.0)			
合計	565	9	574		5	579
(%)	(98.4)	(1.6)	(100.0)			
記載なし	108	2	110		30	140
(%)	(98.2)	(1.8)	(100.0)			
総計	673	11	684		35	719
(%)	(98.4)	(1.6)	(100.0)			
平均	133.22	91.78	132.57		131.20	132.56
標準偏差	130.27	94.26	129.82		71.28	129.39

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査