

表 1459 降圧薬使用の有無 原疾患別（透析患者全体）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	28,579	48,538	77,117	321	19,532	96,970
(%)	(37.1)	(62.9)	(100.0)			
慢性腎盂腎炎	993	1,430	2,423	11	608	3,042
(%)	(41.0)	(59.0)	(100.0)			
急速進行性糸球体腎炎	756	1,236	1,992	9	493	2,494
(%)	(38.0)	(62.0)	(100.0)			
妊娠腎/妊娠中毒症	535	774	1,309	5	300	1,614
(%)	(40.9)	(59.1)	(100.0)			
その他の分類不能の腎炎	370	688	1,058	5	310	1,373
(%)	(35.0)	(65.0)	(100.0)			
多発性嚢胞腎	3,216	5,621	8,837	32	2,137	11,006
(%)	(36.4)	(63.6)	(100.0)			
腎硬化症	7,314	15,627	22,941	113	5,244	28,298
(%)	(31.9)	(68.1)	(100.0)			
悪性高血圧	575	1,477	2,052	5	516	2,573
(%)	(28.0)	(72.0)	(100.0)			
糖尿病性腎症	24,432	69,555	93,987	434	23,660	118,081
(%)	(26.0)	(74.0)	(100.0)			
SLE腎炎	703	1,114	1,817	7	412	2,236
(%)	(38.7)	(61.3)	(100.0)			
アミロイド腎	158	186	344	2	105	451
(%)	(45.9)	(54.1)	(100.0)			
痛風腎	340	581	921	4	188	1,113
(%)	(36.9)	(63.1)	(100.0)			
先天性代謝異常による腎不全	103	114	217	1	54	272
(%)	(47.5)	(52.5)	(100.0)			
腎・尿路結核	86	78	164		42	206
(%)	(52.4)	(47.6)	(100.0)			
腎・尿路結石	184	278	462	1	118	581
(%)	(39.8)	(60.2)	(100.0)			
腎・尿路腫瘍	268	419	687	2	197	886
(%)	(39.0)	(61.0)	(100.0)			
閉塞性尿路障害	263	325	588	1	134	723
(%)	(44.7)	(55.3)	(100.0)			
骨髄腫	105	116	221		55	276
(%)	(47.5)	(52.5)	(100.0)			
腎形成不全	258	278	536	2	111	649
(%)	(48.1)	(51.9)	(100.0)			
不明	7,681	13,584	21,265	144	6,135	27,544
(%)	(36.1)	(63.9)	(100.0)			
再導入	678	1,074	1,752	2	409	2,163
(%)	(38.7)	(61.3)	(100.0)			
その他	2,291	3,542	5,833	41	1,683	7,557
(%)	(39.3)	(60.7)	(100.0)			
合計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			
記載なし						
(%)						
総計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査