

表 1465 降圧薬使用の有無 身長(cm)別（透析患者全体）

身長	なし	あり	合計	不明	記載なし	総計
130未満	148	174	322	2	30	354
(%)	(46.0)	(54.0)	(100.0)			
130～	261	356	617	7	72	696
(%)	(42.3)	(57.7)	(100.0)			
135～	1,017	1,479	2,496	14	327	2,837
(%)	(40.7)	(59.3)	(100.0)			
140～	3,352	5,618	8,970	38	1,289	10,297
(%)	(37.4)	(62.6)	(100.0)			
145～	7,389	13,104	20,493	98	3,082	23,673
(%)	(36.1)	(63.9)	(100.0)			
150～	11,616	21,482	33,098	154	4,983	38,235
(%)	(35.1)	(64.9)	(100.0)			
155～	13,026	25,645	38,671	141	5,660	44,472
(%)	(33.7)	(66.3)	(100.0)			
160～	14,663	31,912	46,575	215	6,860	53,650
(%)	(31.5)	(68.5)	(100.0)			
165～	12,393	29,387	41,780	182	6,026	47,988
(%)	(29.7)	(70.3)	(100.0)			
170～	7,649	19,151	26,800	125	4,149	31,074
(%)	(28.5)	(71.5)	(100.0)			
175～	2,803	7,035	9,838	46	1,435	11,319
(%)	(28.5)	(71.5)	(100.0)			
180～	686	1,903	2,589	14	376	2,979
(%)	(26.5)	(73.5)	(100.0)			
185～	128	280	408	3	54	465
(%)	(31.4)	(68.6)	(100.0)			
190～	19	48	67		10	77
(%)	(28.4)	(71.6)	(100.0)			
合計	75,150	157,574	232,724	1,039	34,353	268,116
(%)	(32.3)	(67.7)	(100.0)			
記載なし	4,738	9,061	13,799	103	28,090	41,992
(%)	(34.3)	(65.7)	(100.0)			
総計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			
平均	159.14	160.47	160.04	159.97	160.08	160.05
標準偏差	9.79	9.57	9.66	10.30	9.58	9.65

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査