

表 1466 降圧薬使用の有無 Body Mass Index別（透析患者全体）

Body Mass Index	なし	あり	合計	不明	記載なし	総計
14未満	650	720	1,370	8	208	1,586
(%)	(47.4)	(52.6)	(100.0)			
14～	3,203	4,870	8,073	50	1,067	9,190
(%)	(39.7)	(60.3)	(100.0)			
16～	9,351	17,373	26,724	110	3,389	30,223
(%)	(35.0)	(65.0)	(100.0)			
18～	15,330	32,308	47,638	188	6,031	53,857
(%)	(32.2)	(67.8)	(100.0)			
20～	16,699	36,415	53,114	202	6,604	59,920
(%)	(31.4)	(68.6)	(100.0)			
22～	12,875	28,136	41,011	164	5,246	46,421
(%)	(31.4)	(68.6)	(100.0)			
24～	7,737	16,988	24,725	97	3,144	27,966
(%)	(31.3)	(68.7)	(100.0)			
26～	4,063	9,133	13,196	60	1,673	14,929
(%)	(30.8)	(69.2)	(100.0)			
28～	2,125	4,730	6,855	21	840	7,716
(%)	(31.0)	(69.0)	(100.0)			
30～	1,781	4,096	5,877	21	713	6,611
(%)	(30.3)	(69.7)	(100.0)			
合計	73,814	154,769	228,583	921	28,915	258,419
(%)	(32.3)	(67.7)	(100.0)			
記載なし	6,074	11,866	17,940	221	33,528	51,689
(%)	(33.9)	(66.1)	(100.0)			
総計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			
平均	21.34	21.60	21.51	21.32	21.48	21.51
標準偏差	3.70	3.63	3.66	3.69	3.66	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査