

表 1473 降圧薬使用の有無 PTH値(pg/mL)別（透析患者全体）PTH測定法:whole-PTH

PTH値	なし	あり	合計	不明	記載なし	総計
20未満	583	1,177	1,760	11	123	1,894
(%)	(33.1)	(66.9)	(100.0)			
20～	715	1,396	2,111	13	207	2,331
(%)	(33.9)	(66.1)	(100.0)			
35～	564	1,313	1,877	10	156	2,043
(%)	(30.0)	(70.0)	(100.0)			
50～	794	2,095	2,889	7	257	3,153
(%)	(27.5)	(72.5)	(100.0)			
75～	655	1,854	2,509	8	231	2,748
(%)	(26.1)	(73.9)	(100.0)			
100～	597	1,476	2,073	7	218	2,298
(%)	(28.8)	(71.2)	(100.0)			
125～	395	963	1,358	5	176	1,539
(%)	(29.1)	(70.9)	(100.0)			
150～	8	34	42		6	48
(%)	(19.0)	(81.0)	(100.0)			
151～	341	787	1,128	4	131	1,263
(%)	(30.2)	(69.8)	(100.0)			
180～	253	524	777	7	91	875
(%)	(32.6)	(67.4)	(100.0)			
210～	178	376	554	5	69	628
(%)	(32.1)	(67.9)	(100.0)			
240～	212	374	586	4	84	674
(%)	(36.2)	(63.8)	(100.0)			
300～	84	163	247	3	34	284
(%)	(34.0)	(66.0)	(100.0)			
350～	60	105	165		14	179
(%)	(36.4)	(63.6)	(100.0)			
400～	62	118	180	3	19	202
(%)	(34.4)	(65.6)	(100.0)			
500～	91	119	210	8	29	247
(%)	(43.3)	(56.7)	(100.0)			
合計	5,592	12,874	18,466	95	1,845	20,406
(%)	(30.3)	(69.7)	(100.0)			
記載なし	139	254	393	4	260	657
(%)	(35.4)	(64.6)	(100.0)			
総計	5,731	13,128	18,859	99	2,105	21,063
(%)	(30.4)	(69.6)	(100.0)			
平均	111.77	104.01	106.36	165.06	121.54	108.01
標準偏差	130.28	100.43	110.38	208.38	138.74	114.01

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査