

表 1474 降圧薬使用の有無 透析前ヘモグロビン濃度(g/dL)別（透析患者全体）

透析前ヘモグロビン濃度	なし	あり	合計	不明	記載なし	総計
5未満	71	207	278		25	303
(%)	(25.5)	(74.5)	(100.0)			
5～	103	279	382		47	429
(%)	(27.0)	(73.0)	(100.0)			
6～	411	575	986	7	167	1,160
(%)	(41.7)	(58.3)	(100.0)			
7～	1,449	2,369	3,818	26	591	4,435
(%)	(38.0)	(62.0)	(100.0)			
8～	4,505	8,159	12,664	68	1,822	14,554
(%)	(35.6)	(64.4)	(100.0)			
9～	12,707	27,261	39,968	176	5,845	45,989
(%)	(31.8)	(68.2)	(100.0)			
10～	25,314	56,192	81,506	316	12,090	93,912
(%)	(31.1)	(68.9)	(100.0)			
11～	21,641	46,346	67,987	276	9,910	78,173
(%)	(31.8)	(68.2)	(100.0)			
12～	8,524	16,648	25,172	124	3,558	28,854
(%)	(33.9)	(66.1)	(100.0)			
13～	2,443	3,910	6,353	27	928	7,308
(%)	(38.5)	(61.5)	(100.0)			
14～	1,021	1,331	2,352	7	314	2,673
(%)	(43.4)	(56.6)	(100.0)			
合計	78,189	163,277	241,466	1,027	35,297	277,790
(%)	(32.4)	(67.6)	(100.0)			
記載なし	1,699	3,358	5,057	115	27,146	32,318
(%)	(33.6)	(66.4)	(100.0)			
総計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			
平均	10.73	10.70	10.71	10.66	10.70	10.71
標準偏差	1.35	1.25	1.29	1.33	1.27	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査