

表 1476 降圧薬使用の有無 HDL-C濃度(mg/dL)別（透析患者全体）

HDL-C濃度	なし	あり	合計	不明	記載なし	総計
20未満	308	389	697	1	98	796
(%)	(44.2)	(55.8)	(100.0)			
20～	4,445	9,293	13,738	28	1,857	15,623
(%)	(32.4)	(67.6)	(100.0)			
30～	12,777	27,519	40,296	87	5,464	45,847
(%)	(31.7)	(68.3)	(100.0)			
40～	14,561	30,618	45,179	107	6,066	51,352
(%)	(32.2)	(67.8)	(100.0)			
50～	10,472	22,434	32,906	74	4,581	37,561
(%)	(31.8)	(68.2)	(100.0)			
60～	6,076	13,400	19,476	48	2,712	22,236
(%)	(31.2)	(68.8)	(100.0)			
70～	3,095	6,449	9,544	28	1,420	10,992
(%)	(32.4)	(67.6)	(100.0)			
80～	1,440	3,066	4,506	5	718	5,229
(%)	(32.0)	(68.0)	(100.0)			
90～	615	1,376	1,991	7	333	2,331
(%)	(30.9)	(69.1)	(100.0)			
100～	537	1,120	1,657		344	2,001
(%)	(32.4)	(67.6)	(100.0)			
合計	54,326	115,664	169,990	385	23,593	193,968
(%)	(32.0)	(68.0)	(100.0)			
記載なし	25,562	50,971	76,533	757	38,850	116,140
(%)	(33.4)	(66.6)	(100.0)			
総計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			
平均	48.55	48.67	48.63	48.70	49.58	48.75
標準偏差	16.65	16.53	16.57	15.34	17.67	16.71

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査