

表 1477 降圧薬使用の有無 Non-HDL-C濃度(mg/dL)別（透析患者全体）

Non-HDL-C濃度	なし	あり	合計	不明	記載なし	総計
50未満	964	1,986	2,950	9	451	3,410
(%)	(32.7)	(67.3)	(100.0)			
50～	4,217	9,769	13,986	42	2,073	16,101
(%)	(30.2)	(69.8)	(100.0)			
70～	9,476	21,421	30,897	93	4,579	35,569
(%)	(30.7)	(69.3)	(100.0)			
90～	11,570	25,365	36,935	90	5,353	42,378
(%)	(31.3)	(68.7)	(100.0)			
110～	9,613	20,316	29,929	62	4,212	34,203
(%)	(32.1)	(67.9)	(100.0)			
130～	6,180	12,124	18,304	33	2,626	20,963
(%)	(33.8)	(66.2)	(100.0)			
150～	3,130	6,140	9,270	18	1,302	10,590
(%)	(33.8)	(66.2)	(100.0)			
170～	1,416	2,732	4,148	3	571	4,722
(%)	(34.1)	(65.9)	(100.0)			
190～	542	1,054	1,596	4	209	1,809
(%)	(34.0)	(66.0)	(100.0)			
210～	324	617	941	3	180	1,124
(%)	(34.4)	(65.6)	(100.0)			
合計	47,432	101,524	148,956	357	21,556	170,869
(%)	(31.8)	(68.2)	(100.0)			
記載なし	32,456	65,111	97,567	785	40,887	139,239
(%)	(33.3)	(66.7)	(100.0)			
総計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			
平均	108.33	106.59	107.15	101.38	106.74	107.08
標準偏差	33.97	33.33	33.54	32.71	34.06	33.61

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査