

表 1478 降圧薬使用の有無 透析前収縮期血圧(mmHg)別（透析患者全体）

透析前収縮期血圧	なし	あり	合計	不明	記載なし	総計
60未満	22	10	32		8	40
(%)	(68.8)	(31.3)	(100.0)			
60～	271	96	367	2	64	433
(%)	(73.8)	(26.2)	(100.0)			
80～	1,980	913	2,893	24	421	3,338
(%)	(68.4)	(31.6)	(100.0)			
100～	8,080	7,257	15,337	80	2,190	17,607
(%)	(52.7)	(47.3)	(100.0)			
120～	17,918	28,850	46,768	181	6,322	53,271
(%)	(38.3)	(61.7)	(100.0)			
140～	21,140	49,912	71,052	304	9,466	80,822
(%)	(29.8)	(70.2)	(100.0)			
160～	13,137	37,945	51,082	188	6,925	58,195
(%)	(25.7)	(74.3)	(100.0)			
180～	4,487	15,022	19,509	72	2,643	22,224
(%)	(23.0)	(77.0)	(100.0)			
200～	1,029	3,865	4,894	16	636	5,546
(%)	(21.0)	(79.0)	(100.0)			
220～	198	819	1,017	2	141	1,160
(%)	(19.5)	(80.5)	(100.0)			
合計	68,262	144,689	212,951	869	28,816	242,636
(%)	(32.1)	(67.9)	(100.0)			
記載なし	11,626	21,946	33,572	273	33,627	67,472
(%)	(34.6)	(65.4)	(100.0)			
総計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			
平均	144.72	154.20	151.16	148.16	150.96	151.13
標準偏差	25.10	23.12	24.18	24.72	24.52	24.23

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査