

表 1479 降圧薬使用の有無 透析前拡張期血圧(mmHg)別（透析患者全体）

透析前拡張 期血圧	なし	あり	合計	不明	記載なし	総計
20未満	26	40	66		14	80
(%)	(39.4)	(60.6)	(100.0)			
20～	395	271	666	7	123	796
(%)	(59.3)	(40.7)	(100.0)			
40～	7,557	10,312	17,869	77	2,534	20,480
(%)	(42.3)	(57.7)	(100.0)			
60～	32,464	65,853	98,317	426	13,135	111,878
(%)	(33.0)	(67.0)	(100.0)			
80～	15,687	36,742	52,429	199	7,059	59,687
(%)	(29.9)	(70.1)	(100.0)			
90～	7,983	20,285	28,268	104	3,879	32,251
(%)	(28.2)	(71.8)	(100.0)			
100～	3,687	10,120	13,807	51	1,913	15,771
(%)	(26.7)	(73.3)	(100.0)			
120～	203	656	859	4	101	964
(%)	(23.6)	(76.4)	(100.0)			
140～	22	41	63		12	75
(%)	(34.9)	(65.1)	(100.0)			
合計	68,024	144,320	212,344	868	28,770	241,982
(%)	(32.0)	(68.0)	(100.0)			
記載なし	11,864	22,315	34,179	274	33,673	68,126
(%)	(34.7)	(65.3)	(100.0)			
総計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			
平均	75.99	78.61	77.77	76.59	77.81	77.77
標準偏差	14.72	14.20	14.42	14.73	14.64	14.45

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査