

表 1480 降圧薬使用の有無 透析前脈拍(拍/分)別 (透析患者全体)

透析前脈拍	なし	あり	合計	不明	記載なし	総計
40未満	63	97	160	1	24	185
(%)	(39.4)	(60.6)	(100.0)			
40～	603	1,513	2,116	7	258	2,381
(%)	(28.5)	(71.5)	(100.0)			
50～	4,817	12,380	17,197	84	2,231	19,512
(%)	(28.0)	(72.0)	(100.0)			
60～	16,237	37,338	53,575	198	7,008	60,781
(%)	(30.3)	(69.7)	(100.0)			
70～	20,208	44,219	64,427	281	8,564	73,272
(%)	(31.4)	(68.6)	(100.0)			
80～	13,779	27,068	40,847	176	5,791	46,814
(%)	(33.7)	(66.3)	(100.0)			
90～	6,447	11,694	18,141	69	2,380	20,590
(%)	(35.5)	(64.5)	(100.0)			
100～	2,233	3,402	5,635	18	740	6,393
(%)	(39.6)	(60.4)	(100.0)			
110～	572	780	1,352	6	170	1,528
(%)	(42.3)	(57.7)	(100.0)			
120～	262	328	590	1	64	655
(%)	(44.4)	(55.6)	(100.0)			
合計	65,221	138,819	204,040	841	27,230	232,111
(%)	(32.0)	(68.0)	(100.0)			
記載なし	14,667	27,816	42,483	301	35,213	77,997
(%)	(34.5)	(65.5)	(100.0)			
総計	79,888	166,635	246,523	1,142	62,443	310,108
(%)	(32.4)	(67.6)	(100.0)			
平均	75.68	74.19	74.66	74.52	74.88	74.69
標準偏差	13.24	12.66	12.86	12.50	12.64	12.84

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査