

表 1484 降圧薬使用の有無 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	21,581	36,951	58,532	261	9,784	68,577
(%)	(36.9)	(63.1)	(100.0)			
慢性腎盂腎炎	751	1,100	1,851	7	307	2,165
(%)	(40.6)	(59.4)	(100.0)			
急速進行性糸球体腎炎	625	977	1,602	9	223	1,834
(%)	(39.0)	(61.0)	(100.0)			
妊娠腎/妊娠中毒症	362	554	916	3	145	1,064
(%)	(39.5)	(60.5)	(100.0)			
その他の分類不能の腎炎	263	519	782	3	164	949
(%)	(33.6)	(66.4)	(100.0)			
多発性嚢胞腎	2,569	4,485	7,054	26	1,118	8,198
(%)	(36.4)	(63.6)	(100.0)			
腎硬化症	6,078	12,608	18,686	86	2,688	21,460
(%)	(32.5)	(67.5)	(100.0)			
悪性高血圧	476	1,144	1,620	4	257	1,881
(%)	(29.4)	(70.6)	(100.0)			
糖尿病性腎症	20,310	57,305	77,615	362	12,530	90,507
(%)	(26.2)	(73.8)	(100.0)			
SLE腎炎	532	832	1,364	4	182	1,550
(%)	(39.0)	(61.0)	(100.0)			
アミロイド腎	129	141	270	2	53	325
(%)	(47.8)	(52.2)	(100.0)			
痛風腎	274	455	729	2	97	828
(%)	(37.6)	(62.4)	(100.0)			
先天性代謝異常による腎不全	76	80	156		21	177
(%)	(48.7)	(51.3)	(100.0)			
腎・尿路結核	68	71	139		19	158
(%)	(48.9)	(51.1)	(100.0)			
腎・尿路結石	155	214	369	1	60	430
(%)	(42.0)	(58.0)	(100.0)			
腎・尿路腫瘍	227	350	577	1	108	686
(%)	(39.3)	(60.7)	(100.0)			
閉塞性尿路障害	197	263	460	1	56	517
(%)	(42.8)	(57.2)	(100.0)			
骨髄腫	80	91	171		19	190
(%)	(46.8)	(53.2)	(100.0)			
腎形成不全	174	200	374		45	419
(%)	(46.5)	(53.5)	(100.0)			
不明	6,137	10,732	16,869	118	3,138	20,125
(%)	(36.4)	(63.6)	(100.0)			
再導入	484	756	1,240		177	1,417
(%)	(39.0)	(61.0)	(100.0)			
その他	1,818	2,808	4,626	33	855	5,514
(%)	(39.3)	(60.7)	(100.0)			
合計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			
記載なし						
(%)						
総計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査