

表 1489 降圧薬使用の有無 透析時間(時間/回)別(施設血液透析患者 透析回数週3回)

透析時間	なし	あり	合計	不明	記載なし	総計
3.0未満	183	261	444	6	71	521
(%)	(41.2)	(58.8)	(100.0)			
3.0～	6,932	14,053	20,985	145	3,690	24,820
(%)	(33.0)	(67.0)	(100.0)			
3.5～	4,724	10,636	15,360	78	2,501	17,939
(%)	(30.8)	(69.2)	(100.0)			
4.0～	42,320	90,643	132,963	626	21,409	154,998
(%)	(31.8)	(68.2)	(100.0)			
4.5～	3,256	7,045	10,301	42	1,335	11,678
(%)	(31.6)	(68.4)	(100.0)			
5.0～	4,944	8,787	13,731	22	1,834	15,587
(%)	(36.0)	(64.0)	(100.0)			
5.5～	230	317	547		48	595
(%)	(42.0)	(58.0)	(100.0)			
6.0～	410	480	890	1	103	994
(%)	(46.1)	(53.9)	(100.0)			
6.5～	29	28	57		1	58
(%)	(50.9)	(49.1)	(100.0)			
7.0～	236	217	453		58	511
(%)	(52.1)	(47.9)	(100.0)			
合計	63,264	132,467	195,731	920	31,050	227,701
(%)	(32.3)	(67.7)	(100.0)			
記載なし	102	169	271	3	996	1,270
(%)	(37.6)	(62.4)	(100.0)			
総計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			
平均	3.99	3.97	3.98	3.85	3.95	3.97
標準偏差	0.56	0.51	0.53	0.46	0.51	0.52

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査