

表 1491 降圧薬使用の有無 身長(cm)別（施設血液透析患者 透析回数週3回）

身長	なし	あり	合計	不明	記載なし	総計
130未満	103	113	216	2	19	237
(%)	(47.7)	(52.3)	(100.0)			
130～	218	291	509	6	53	568
(%)	(42.8)	(57.2)	(100.0)			
135～	820	1,219	2,039	12	254	2,305
(%)	(40.2)	(59.8)	(100.0)			
140～	2,742	4,576	7,318	26	939	8,283
(%)	(37.5)	(62.5)	(100.0)			
145～	5,994	10,641	16,635	81	2,103	18,819
(%)	(36.0)	(64.0)	(100.0)			
150～	9,223	17,165	26,388	132	3,494	30,014
(%)	(35.0)	(65.0)	(100.0)			
155～	10,277	20,524	30,801	118	3,938	34,857
(%)	(33.4)	(66.6)	(100.0)			
160～	11,671	25,590	37,261	181	4,816	42,258
(%)	(31.3)	(68.7)	(100.0)			
165～	9,847	23,387	33,234	141	4,201	37,576
(%)	(29.6)	(70.4)	(100.0)			
170～	5,886	14,971	20,857	101	2,869	23,827
(%)	(28.2)	(71.8)	(100.0)			
175～	2,128	5,434	7,562	36	951	8,549
(%)	(28.1)	(71.9)	(100.0)			
180～	538	1,449	1,987	12	255	2,254
(%)	(27.1)	(72.9)	(100.0)			
185～	105	210	315	3	33	351
(%)	(33.3)	(66.7)	(100.0)			
190～	14	38	52		5	57
(%)	(26.9)	(73.1)	(100.0)			
合計	59,566	125,608	185,174	851	23,930	209,955
(%)	(32.2)	(67.8)	(100.0)			
記載なし	3,800	7,028	10,828	72	8,116	19,016
(%)	(35.1)	(64.9)	(100.0)			
総計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			
平均	159.04	160.36	159.94	159.86	159.99	159.94
標準偏差	9.71	9.51	9.59	10.31	9.54	9.59

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査