

表 1492 降圧薬使用の有無 透析後体重(kg)別（施設血液透析患者 透析回数週3回）

透析後体重	なし	あり	合計	不明	記載なし	総計
20未満	21	54	75	1	8	84
(%)	(28.0)	(72.0)	(100.0)			
20～	323	358	681	5	116	802
(%)	(47.4)	(52.6)	(100.0)			
30～	6,500	10,081	16,581	81	2,455	19,117
(%)	(39.2)	(60.8)	(100.0)			
40～	17,854	34,147	52,001	230	7,578	59,809
(%)	(34.3)	(65.7)	(100.0)			
50～	19,649	43,074	62,723	290	9,096	72,109
(%)	(31.3)	(68.7)	(100.0)			
60～	11,544	26,829	38,373	163	5,546	44,082
(%)	(30.1)	(69.9)	(100.0)			
70～	4,431	10,547	14,978	62	2,144	17,184
(%)	(29.6)	(70.4)	(100.0)			
80～	1,538	3,776	5,314	21	746	6,081
(%)	(28.9)	(71.1)	(100.0)			
90～	532	1,383	1,915	9	271	2,195
(%)	(27.8)	(72.2)	(100.0)			
100～	170	482	652	4	99	755
(%)	(26.1)	(73.9)	(100.0)			
110～	56	201	257	2	40	299
(%)	(21.8)	(78.2)	(100.0)			
120～	49	110	159	1	29	189
(%)	(30.8)	(69.2)	(100.0)			
合計	62,667	131,042	193,709	869	28,128	222,706
(%)	(32.4)	(67.6)	(100.0)			
記載なし	699	1,594	2,293	54	3,918	6,265
(%)	(30.5)	(69.5)	(100.0)			
総計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			
平均	54.41	55.97	55.46	54.93	55.34	55.45
標準偏差	12.81	12.90	12.89	13.23	12.91	12.90

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査