

表 1495 降圧薬使用の有無 Body Mass Index別（施設血液透析患者 透析回数週3回）

Body Mass Index	なし	あり	合計	不明	記載なし	総計
14未満	540	594	1,134	7	170	1,311
(%)	(47.6)	(52.4)	(100.0)			
14～	2,664	4,076	6,740	46	820	7,606
(%)	(39.5)	(60.5)	(100.0)			
16～	7,601	14,159	21,760	97	2,623	24,480
(%)	(34.9)	(65.1)	(100.0)			
18～	12,141	26,129	38,270	163	4,723	43,156
(%)	(31.7)	(68.3)	(100.0)			
20～	13,111	29,276	42,387	171	5,122	47,680
(%)	(30.9)	(69.1)	(100.0)			
22～	10,212	22,398	32,610	143	4,057	36,810
(%)	(31.3)	(68.7)	(100.0)			
24～	6,183	13,218	19,401	80	2,432	21,913
(%)	(31.9)	(68.1)	(100.0)			
26～	3,205	7,122	10,327	48	1,313	11,688
(%)	(31.0)	(69.0)	(100.0)			
28～	1,676	3,646	5,322	18	651	5,991
(%)	(31.5)	(68.5)	(100.0)			
30～	1,412	3,187	4,599	17	548	5,164
(%)	(30.7)	(69.3)	(100.0)			
合計	58,745	123,805	182,550	790	22,459	205,799
(%)	(32.2)	(67.8)	(100.0)			
記載なし	4,621	8,831	13,452	133	9,587	23,172
(%)	(34.4)	(65.6)	(100.0)			
総計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			
平均	21.31	21.53	21.46	21.23	21.47	21.46
標準偏差	3.72	3.62	3.66	3.69	3.66	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査