

表 1496 降圧薬使用の有無 透析前BUN(mg/dL)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前BUN	なし	あり	合計	不明	記載なし	総計
20未満	178	228	406	2	71	479
(%)	(43.8)	(56.2)	(100.0)			
20～	850	1,306	2,156	12	312	2,480
(%)	(39.4)	(60.6)	(100.0)			
30～	3,227	5,721	8,948	43	1,367	10,358
(%)	(36.1)	(63.9)	(100.0)			
40～	7,921	15,387	23,308	86	3,417	26,811
(%)	(34.0)	(66.0)	(100.0)			
50～	12,068	24,933	37,001	147	5,511	42,659
(%)	(32.6)	(67.4)	(100.0)			
60～	11,949	24,969	36,918	148	5,577	42,643
(%)	(32.4)	(67.6)	(100.0)			
70～	7,985	16,650	24,635	116	3,571	28,322
(%)	(32.4)	(67.6)	(100.0)			
80～	3,919	7,567	11,486	59	1,667	13,212
(%)	(34.1)	(65.9)	(100.0)			
90～	1,412	2,747	4,159	15	629	4,803
(%)	(34.0)	(66.0)	(100.0)			
100～	413	770	1,183	10	155	1,348
(%)	(34.9)	(65.1)	(100.0)			
110～	109	238	347		47	394
(%)	(31.4)	(68.6)	(100.0)			
120～	70	103	173	1	30	204
(%)	(40.5)	(59.5)	(100.0)			
合計	50,101	100,619	150,720	639	22,354	173,713
(%)	(33.2)	(66.8)	(100.0)			
記載なし	423	1,056	1,479	39	2,978	4,496
(%)	(28.6)	(71.4)	(100.0)			
総計	50,524	101,675	152,199	678	25,332	178,209
(%)	(33.2)	(66.8)	(100.0)			
平均	60.69	61.08	60.95	61.71	60.84	60.94
標準偏差	16.05	15.43	15.64	16.33	15.62	15.64

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査