

表 1505 降圧薬使用の有無 透析前リン濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

透析前リン濃度	なし	あり	合計	不明	記載なし	総計
3未満	3,186	4,553	7,739	36	1,186	8,961
(%)	(41.2)	(58.8)	(100.0)			
3.0～	3,399	6,104	9,503	37	1,440	10,980
(%)	(35.8)	(64.2)	(100.0)			
3.5～	5,598	11,015	16,613	74	2,331	19,018
(%)	(33.7)	(66.3)	(100.0)			
4.0～	8,090	16,538	24,628	113	3,451	28,192
(%)	(32.8)	(67.2)	(100.0)			
4.5～	9,380	19,869	29,249	123	4,096	33,468
(%)	(32.1)	(67.9)	(100.0)			
5.0～	9,150	20,239	29,389	123	4,297	33,809
(%)	(31.1)	(68.9)	(100.0)			
5.5～	8,019	17,334	25,353	102	3,557	29,012
(%)	(31.6)	(68.4)	(100.0)			
6.0～	1,346	2,956	4,302	9	617	4,928
(%)	(31.3)	(68.7)	(100.0)			
6.1～	4,419	9,922	14,341	69	2,095	16,505
(%)	(30.8)	(69.2)	(100.0)			
6.5～	3,810	8,698	12,508	65	1,804	14,377
(%)	(30.5)	(69.5)	(100.0)			
7.0～	2,334	5,363	7,697	37	1,126	8,860
(%)	(30.3)	(69.7)	(100.0)			
7.5～	1,451	3,240	4,691	24	644	5,359
(%)	(30.9)	(69.1)	(100.0)			
8.0～	853	1,951	2,804	15	373	3,192
(%)	(30.4)	(69.6)	(100.0)			
8.5～	575	1,150	1,725	13	285	2,023
(%)	(33.3)	(66.7)	(100.0)			
9.0～	870	1,886	2,756	17	383	3,156
(%)	(31.6)	(68.4)	(100.0)			
合計	62,480	130,818	193,298	857	27,685	221,840
(%)	(32.3)	(67.7)	(100.0)			
記載なし	886	1,818	2,704	66	4,361	7,131
(%)	(32.8)	(67.2)	(100.0)			
総計	63,366	132,636	196,002	923	32,046	228,971
(%)	(32.3)	(67.7)	(100.0)			
平均	5.12	5.23	5.20	5.29	5.19	5.20
標準偏差	1.45	1.41	1.42	1.54	1.44	1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査