

表 1507 降圧薬使用の有無 PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	なし	あり	合計	不明	記載なし	総計
20未満	457	946	1,403	9	87	1,499
(%)	(32.6)	(67.4)	(100.0)			
20～	592	1,122	1,714	11	123	1,848
(%)	(34.5)	(65.5)	(100.0)			
35～	447	1,050	1,497	9	107	1,613
(%)	(29.9)	(70.1)	(100.0)			
50～	603	1,689	2,292	7	160	2,459
(%)	(26.3)	(73.7)	(100.0)			
75～	508	1,481	1,989	6	153	2,148
(%)	(25.5)	(74.5)	(100.0)			
100～	479	1,161	1,640	7	140	1,787
(%)	(29.2)	(70.8)	(100.0)			
125～	308	756	1,064	4	119	1,187
(%)	(28.9)	(71.1)	(100.0)			
150～	7	26	33		4	37
(%)	(21.2)	(78.8)	(100.0)			
151～	255	598	853	4	77	934
(%)	(29.9)	(70.1)	(100.0)			
180～	194	408	602	7	57	666
(%)	(32.2)	(67.8)	(100.0)			
210～	132	283	415	4	53	472
(%)	(31.8)	(68.2)	(100.0)			
240～	173	295	468	4	54	526
(%)	(37.0)	(63.0)	(100.0)			
300～	72	111	183	3	19	205
(%)	(39.3)	(60.7)	(100.0)			
350～	46	78	124		11	135
(%)	(37.1)	(62.9)	(100.0)			
400～	52	89	141	3	8	152
(%)	(36.9)	(63.1)	(100.0)			
500～	68	80	148	8	10	166
(%)	(45.9)	(54.1)	(100.0)			
合計	4,393	10,173	14,566	86	1,182	15,834
(%)	(30.2)	(69.8)	(100.0)			
記載なし	76	177	253	3	182	438
(%)	(30.0)	(70.0)	(100.0)			
総計	4,469	10,350	14,819	89	1,364	16,272
(%)	(30.2)	(69.8)	(100.0)			
平均	111.36	101.75	104.65	174.76	113.63	105.70
標準偏差	131.50	97.95	109.24	215.63	95.04	109.24

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査